
My NASA Data - Lesson Plans

Villarrica Volcano Eruption

Grade Band

- 6-8

Time

- 50 minutes

Overview

Students use scale to determine the area of volcanic deposits following the March 3, 2015 eruption of Chile's Mount Villarrica stratovolcano, one of the country's most active volcanoes.

Materials Required

- Rulers
- Colored printouts of images

Procedure

Introduce the Event:

1. Show students the two pictures of Mount Villarrica. Do not tell them what they are. Have them make observations about the pictures and note similarities and differences.

Work Collaboratively: Think-Pair-Share

1. Have students discuss and share with each other what they think happened in the picture.
2. After students share their ideas, provide the context for the pictures.

In the predawn hours of March 3, 2015, one of Chile's most active volcanoes exploded to life. Thousands of people evacuated from within a 10-kilometer (6-mile) radius of Villarrica.



Villarrica Volcano February 22, 2015- Image NASA Earth Observatory

The Operational Land Imager (OLI) on Landsat 8 acquired the top image on February 22, 2015, *prior to the main eruption*. The Advanced Land Imager (ALI) on the EO-1 satellite captured the second image on March 5. A large volume of volcanic material ejected during the eruption is apparent on the mountain's eastern side.

Activity at Villarrica had been increasing through much of February, made evident by a rise in seismicity, crater incandescence, and explosions, according to reports from Chile's National Geology and Mining Service (SERNAGEOMIN) (posted in English by the Smithsonian Institution's Global Volcanism Program). A faint layer of ash is apparent on the volcano's flanks.

Just prior to the March 3 eruption, geophysicist Jeffrey Johnson and his family prepared to evacuate their home in Pucón, Chile, about 17 kilometers (11 miles) northwest of the volcano. However, the eruption slowed after just 30 minutes. Johnson watched the fire fountain subside and then resumed his work analyzing low-frequency sounds captured by dozens of microphones placed on and near the mountain.

NASA Earth Observatory images by Jesse Allen, using EO-1 ALI data provided courtesy of the NASA EO-1 team and Landsat data from the U.S. Geological Survey. Caption by Kathryn Hansen.




Villarrica Volcano March 5, 2015- Image Credit NASA Earth Observatory

Determine the Affected Area:

Students may work in small groups or individually to calculate the area covered by volcanic debris.

1. Direct students to focus on the March 5, 2015 image.
2. Ask students to estimate the area of the new volcanic deposits.
3. Measure the length on the image scale that is equal to 1 km. Call the measurement S.
4. Have students write the scale $S \text{ cm} = 1 \text{ km}$.
5. Draw a rectangle around the area debris that was deposited. It will not fit exactly but should be their best estimate.



Sample rectangle for area calculation

$$\frac{1 \text{ cm}}{1 \text{ km}} = \frac{3.6 \text{ cm}}{x \text{ km}}$$

$$(1 \text{ cm})(x \text{ km}) = (3.6 \text{ cm})(1 \text{ km})$$

$$x \text{ km} = \frac{(3.6 \text{ cm})(1 \text{ km})}{1 \text{ cm}}$$

$$x \text{ km} = \frac{(3.6 \cancel{\text{cm}})(1 \text{ km})}{1 \cancel{\text{cm}}}$$

$$x \text{ km} = 3.6 \text{ km}$$

Sample calculation for length

using scale

6. Measure the length and width of the rectangle.
7. Use the scale on the image to calculate the area of the debris in square kilometers.
8. Example (For illustration purposes only. Assume the scale is 1 cm = 1 km and the length and width are the measurements below.):
 - Length = 3.2 cm
 - Width = 1.6 cm

-
- Calculate the length as follows.
9. Calculate the width the same way.
 10. Calculate the area of the area by multiplying the length by the width. Remember to use the correct units.
 11. The eruption lasted approximately 30 minutes. Calculate the rate of deposition by dividing the area by the time.
 - Example (For illustration purposes only. Assume the area is 25 km²):

$$\frac{25 \text{ km}^2}{30 \text{ min}} = \frac{0.83 \text{ km}^2}{\text{min}}$$

Sample rate calculation.

Exit Ticket

- Based upon the calculations, what would be a safe distance from the volcano if there were an hour long eruption of the same force? Explain.

NGSS Three Dimensional Learning

NGSS Disciplinary Core Ideas

- ESS2B: Plate Tectonics and Large-Scale Systems

NGSS Crosscutting Concepts

- Scale, Proportion, and Quantity

NGSS Science and Engineering Practices

- Developing and Using Models

Supported Common Core Math

- CCSS.Math.Content.6.RP.A.3.B
- CCSS.Math.Content.6.RP.A.3.D
- CCSS.Math.Content.7.RP.A.1

Learning Objectives

- Students use a scale to calculate a map area.
- Students determine the area of volcanic deposits.
- Students explain the time scale involved in the measured change.

Essential Questions

- How quickly did volcanic debris from Mount Villarricia's eruption cover the area?
- How much of the area was covered?

Why Does NASA Study This Phenomenon?

Landsat 8

Launched on February 11, 2013, Landsat 8 (formerly the Landsat Data Continuity Mission, LDCM) is the most recently launched Landsat satellite. It is collecting valuable data and imagery used in agriculture, education, business, science, and government.

The Landsat Program provides repetitive acquisition of high resolution multispectral data of the Earth's surface on a global basis. The data from Landsat spacecraft constitute the longest record of the Earth's continental surfaces as seen from space. It is a record unmatched in quality, detail, coverage, and value.

[Video: Landsat Data Continuity Mission Overview](https://www.youtube.com/watch?v=mqVKR9OnqqA)

Landsat Data Continuity Mission Overview | <https://www.youtube.com/watch?v=mqVKR9OnqqA> |
Source: NASA Video

Teacher Background Information

Glacier-clad Villarrica is one of Chile's most active volcanoes. It rises above the lake and town of the same name. It is the westernmost of three large stratovolcanoes that trend perpendicular to the Andean chain. A 6-km-wide caldera formed during the late Pleistocene. A 2-km-wide caldera that formed about 3500 years ago is located at the base of the presently active, dominantly basaltic to basaltic-andesitic cone at the NW margin of the Pleistocene caldera. Historical eruptions, documented since 1558, have consisted largely of mild-to-moderate explosive activity with occasional lava effusion.

“Global Volcanism Program: Villarrica.” *Global Volcanism Program | Villarrica*, Global Volcanism Program: Villarrica, <https://volcano.si.edu/volcano.cfm?vn=357120>.

For additional background information on volcanoes, see the [Volcanic Eruptions](#) phenomenon page.

Student Misconception

- Volcanoes are randomly located across the earth's surface.
- Volcanoes are found only on land.
- Volcanoes are found only in hot climates.
- All volcanoes erupt violently.
- Volcanoes only erupt straight up through the top vent.
- If a volcano doesn't erupt for a hundred years, it's extinct.
- If a volcano does not produce lava, it is not dangerous.

Source: [Common Misconceptions about Weathering, Erosion, Volcanoes, and Earthquakes](#)

Student Resources

- [Villarrica Volcano Eruption Images](#)